

Plasminogen Deficiency, Type 1: For ENT/Otolaryngology



Overview of Plasminogen Deficiency (PLGD)

- PLGD Type 1: a quantitative protein deficiency, with decreased plasminogen activity and decreased antigen
 - Due to a genetic mutation in the gene for plasminogen; > 50 different mutations have been identified; Autosomal Recessive inheritance
 - Most common presenting symptom is ligneous conjunctivitis, but multi-organ, systemic disease that can be life-threatening
- PLGD Type 2: reduced functional activity of plasminogen, but normal antigen levels; patients are asymptomatic


	Normal	PLGD Type 1	PLGD Type 2
Plasminogen Activity	70-130%	Decreased	Decreased
Plasminogen Antigen	6-25 mg/dL	Decreased	Normal
For Patients: My Plasminogen Activity			

Diagnosis

- Complicated by heterogeneous symptoms; symptoms can wax and wane
- Mucosal surfaces of the eyes, ears, nose, gums, airways, lungs, GI tract, kidneys, GU tract, CNS, and skin can all be affected
- Initial point of medical contact therefore includes many disciplines


Treatment

- Ryplazim (plasminogen, human-tvmh) given by IV infusion leads to resolution of lesions
- Surgical removal of lesions, though initially helpful, leads to accelerated regrowth
- Referral to a Hemophilia Treatment Center (HTC) to serve as medical home, and:
 - Educate on product use
 - Ongoing symptom monitoring, outcomes, safety
 - Administer doses, determine dosing schedule
 - Teach home infusion



*Lesion removal prior to diagnosis not recommended

ENT / Otolaryngology Specific Diagnostic Findings



ENT/Otolaryngology Specific Treatment Considerations

- New or suspected diagnosis:
 - Obtain diagnostic blood test (plasminogen activity level) or refer to hematologist to order
 - If confirmed, refer to HTC to establish care and for thorough review of systems
 - Patients may have more than one system affected at presentation or occurring over time
 - Early referral to HTC is critical to reduce risk of permanent sequelae
- Confirmed diagnosis:
 - Coordinate clinical care with existing care team at HTC; send clinical notes and photos and work closely in collaboration with HTC
- Be suspicious of common diagnoses in PLGD patients, as they may be the result of the condition
 - Chronic hoarseness due to edema or lesions of the vocal cords
 - Chronic sinusitis
 - Nasopharyngeal involvement
 - Nasal cavity lesions leading to irritation, epistaxis, and obstruction
 - Recurrent tonsillitis, with or without tonsilloliths
 - Recurrent ear infections with frequent malfunction of myringotomy tubes (e.g., otorrhea, tube obstruction/extrusion), or otitis media with effusion (OME)
 - Laryngotracheobronchial tree involvement
 - Cholesteatoma-like complications of the middle ear with involvement of the mastoid system
 - Ligneous conjunctivitis (often the first noticeable symptom of PLGD)
- Coordination with HTC assures optimal outcomes from interventions (e.g., bronchoscopy, laryngoscopy, PE tube placement)

Learn More:



Partners in
Bleeding Disorders
Education



PLASMINOGEN
DEFICIENCY
FOUNDATION

